arXiv:0809.2865v1 [math-ph] 17 Sep 2008

About the seventh-order Kaup-Kupershmidt equation and its solutions

Alvaro H. Salas S.
email : asalash2002@yahoo.com
Department of Mathematics
Universidad de Caldas, Manizales, Colombia.
Universidad Nacional de Colombia, Manizales.

Jairo Ernesto Castillo Hernández
Department of Physics
Universidad Autónoma, Bogotá, Colombia.
Universidad Central, Bogotá, Colombia.

José Gonzalo Escobar Lugo
email : jogoel@gmail.com
Universidad Cooperativa de Colombia, Bogotá.
Department of Mathematics
Universidad Libre, Bogotá, Colombia.

Abstract. In this letter we obtain exact soliton and periodic solutions to the seventh-order Kaup-Kupershmidt equation. We make use of the Cole-Hopf transformation and two particular rational hyperbolic functions ansatze.

Keywords and phrases. Nonlinear differential equation, nonlinear partial differential equation, seventh-order evolution equation, KdV7, soliton solution, Kaup-Kupershmidt equation.

2000 Mathematics Subject Classification. 35C05.

1. Introduction

The general form of the seventh order KdV (KdV7) [2] reads

$$u_t + au^3u_x + bu_x^3 + cuu_xu_{xx} + du^2u_{xxx} + eu_{2x}u_{3x} + fu_xu_{4x} + guu_{5x} + u_{7x} = 0. \tag{1.1}$$

The seventh-order KdV equation has been introduced by Pomeau et al. [1] for discussing the structural stability of KdV equation under a singular perturbation. Some particular cases of Eq. (1.1) are :

- Seventh-order Sawada-Kotera-Ito equation [2] ($a = 252$, $b = 63$, $c = 378$, $d = 126$, $e = 63$, $f = 42$, $g = 21$) :

$$u_t + 252u^3u_x + 63u_x^3 + 378uu_xu_{xx} + 126u^2u_{xxx} + 63u_{2x}u_{3x} + 42u_xu_{4x} + 21uu_{5x} + u_{7x} = 0. \tag{1.2}$$

- Seventh-order Lax equation [2] ($a = 140$, $b = 70$, $c = 280$, $d = 70$, $e = 70$, $f = 42$, $g = 14$) :

$$u_t + 140u^3u_x + 70u_x^3 + 280uu_xu_{xx} + 70u^2u_{xxx} + 70u_{2x}u_{3x} + 42u_xu_{4x} + 14uu_{5x} + u_{7x} = 0. \tag{1.3}$$

- Seventh-order Kaup-Kupershmidt equation [2] ($a = 2016$, $b = 630$, $c = 2268$, $d = 504$, $e = 252$, $f = 147$, $g = 42$) :

$$u_t + 2016u^3u_x + 630u_x^3 + 2268uu_xu_{xx} + 504u^2u_{xxx} + 252u_{2x}u_{3x} + 147u_xu_{4x} + 42uu_{5x} + u_{7x} = 0. \tag{1.4}$$

In the next sections we obtain some exact solutions for the seventh-order Kaup-Kupershmidt equation (1.4).

2. Soliton solutions by the Cole-Hopf transformation

Some nonlinear pse's in the variable $u = u(x, t)$ may be solved via the substitution

$$u(x, t) = A\partial_{xx} \log(1 + f(x, t)) + B, \tag{2.1}$$

where A and B are some constants and $f(x, t)$ is a new unknown function. Eq. (2.1) is called a generalized Cole-Hopf transformation [3][4]. We will solve Eq. (1.4) by using (2.1) for the special choice

$$f(x, t) = \exp(kx - ct + \delta),$$

being c, k and δ some constants. Now, we make the substitution

$$u(x, t) = A\partial_{xx} \log(1 + \exp(kx - ct + \delta)) + B \tag{2.2}$$

into (1.4). We obtain a polynomial equation in the variable $\zeta = \exp(kx - ct + \delta)$. Equating the coefficients of the different powers of ζ to zero, we obtain the following algebraic system :

- $-Ak^9 - 42ABk^7 - 504AB^2k^5 - 2016AB^3k^3 + Ack^2 = 0.$
- $Ak^9 + 42ABk^7 + 504AB^2k^5 + 2016AB^3k^3 - Ack^2 = 0.$
- $-441A^2k^9 + 247Ak^9 - 3276A^2Bk^7 + 2310ABk^7 - 6048A^2B^2k^5 + 3528AB^2k^5 - 10080AB^3k^3 + 5Ack^2 = 0.$
- $441A^2k^9 - 247Ak^9 + 3276A^2Bk^7 - 2310ABk^7 + 6048A^2B^2k^5 - 3528AB^2k^5 + 10080AB^3k^3 - 5Ack^2 = 0.$
- $-3402A^3k^9 + 10143A^2k^9 - 4293Ak^9 - 6048A^3Bk^7 + 15876A^2Bk^7 - 7938ABk^7 - 18144A^2B^2k^5 + 13608AB^2k^5 - 18144AB^3k^3 + 9Ack^2 = 0.$
- $3402A^3k^9 - 10143A^2k^9 + 4293Ak^9 + 6048A^3Bk^7 - 15876A^2Bk^7 + 7938ABk^7 + 18144A^2B^2k^5 - 13608AB^2k^5 + 18144AB^3k^3 - 9Ack^2 = 0.$
- $-2016A^4k^9 + 18774A^3k^9 - 40320A^2k^9 + 15619Ak^9 - 6048A^3Bk^7 + 19152A^2Bk^7 - 10290ABk^7 - 12096A^2B^2k^5 + 9576AB^2k^5 - 10080AB^3k^3 + 5Ack^2 = 0.$
- $2016A^4k^9 - 18774A^3k^9 + 40320A^2k^9 - 15619Ak^9 + 6048A^3Bk^7 - 19152A^2Bk^7 + 10290ABk^7 + 12096A^2B^2k^5 - 9576AB^2k^5 + 10080AB^3k^3 - 5Ack^2 = 0.$

Solving this systems gives

$$A = \frac{1}{2}, \qquad B = -\frac{k^2}{24}, \qquad c = -\frac{k^7}{48}. \tag{2.3}$$

Therefore, a one-soliton solution of (1.4) is given by

$$u_0(x, t) = -\frac{\left(1 - 10e^{\frac{tk^7}{48} + xk + \delta} + e^{\frac{tk^7}{24} + 2xk + 2\delta}\right)k^2}{24\left(1 + e^{\frac{tk^7}{48} + xk + \delta}\right)^2}, \tag{2.4}$$

which simplifies to

$$u_0(x,t) = -\frac{k^2}{24} + \frac{1}{4(1+\cosh\left(kx + \frac{k^7}{48}t + \delta\right))}. \tag{2.5}$$

3. Exact solutions via two rational hyperbolic ansatze

Using the wave transformation

$$u(x,t) = v(\xi),\ \xi = x + \lambda t, \tag{3.1}$$

where λ is a constant, Eq. (1.4) becomes the nonlinear ode

$$\begin{aligned}&2016v'(\xi)u(\xi)^3 + 504v^{(3)}(\xi)v(\xi)^2 + 2268v'(\xi)v''(\xi)v(\xi) + 42v^{(5)}(\xi)v(\xi) + 630v'(\xi)^3 + \\ &\lambda v'(\xi) + 252v''(\xi)v^{(3)}(\xi) + 147v'(\xi)v^{(4)}(\xi) + v^{(7)}(\xi) = 0.\end{aligned} \tag{3.2}$$

We shall find solutions of Eq. (3.2) by using two ansatze:

- **The tanh–coth ansatz**:

$$v(\xi) = p + a\tanh(\mu\xi) + b\coth(\mu\xi) + c\tanh^2(\mu\xi) + d\coth^2(\mu\xi) \tag{3.3}$$

- **A sinh–cosh rational ansatz**:

$$v(\xi) = p + \frac{k}{1 + c\sinh(\mu\xi) + d\cosh(\mu\xi)}, \tag{3.4}$$

where a, b, c, d, k, p and μ are constants. Some times, by replacing μ by $\sqrt{-1}\mu$, we obtain periodic solutions.

3.1. Solutions by the tanh–coth ansatz.

We change the tanh and coth functions to their exponential form and then we substitute (3.3) into (3.2). We obtain a polynomial equation in the variable $\zeta = \exp(\mu\xi)$. Equating the coefficients of the different powers of ζ to zero results in an algebraic system in the variables a, b, c, d, p, λ and μ. Solving it with the aid of a computer, we obtain following solutions of (1.4) :

- $a = 0, \quad b = 0, \quad c = 0, \quad d = -\frac{\mu^2}{2}, \quad p = \frac{\mu^2}{3}, \quad \lambda = \frac{4\mu^6}{3}$:

$$u_1(x,t) = \frac{\mu^2}{3} - \frac{\mu^2}{2}\coth^2\left(\mu\left(x + \frac{4\mu^6}{3}t\right)\right).$$

$$u_2(x,t) = -\frac{\mu^2}{3} - \frac{\mu^2}{2}\cot^2\left(\mu\left(x - \frac{4\mu^6}{3}t\right)\right).$$

- $a = 0, \quad b = 0, \quad c = -\frac{\mu^2}{2}, \quad d = 0, \quad p = \frac{\mu^2}{3}, \quad \lambda = \frac{4\mu^6}{3}$:

$$u_3(x,t) = \frac{\mu^2}{3} - \frac{\mu^2}{2}\tanh^2\left(\mu\left(x + \frac{4\mu^6}{3}t\right)\right).$$

$$u_4(x,t) = -\frac{\mu^2}{3} - \frac{\mu^2}{2}\tan^2\left(\mu\left(x - \frac{4\mu^6}{3}t\right)\right).$$

- $a = 0, \quad b = 0, \quad c = -\frac{\mu^2}{2}, \quad d = -\frac{\mu^2}{2}, \quad p = \frac{\mu^2}{3}, \quad \lambda = \frac{256\mu^6}{3}$:

$$u_5(x,t) = \frac{\mu^2}{3} - \frac{\mu^2}{2}\coth^2\left(\mu\left(x + \frac{256\mu^6}{3}t\right)\right) - \frac{\mu^2}{2}\tanh^2\left(\mu\left(x + \frac{256\mu^6}{3}t\right)\right).$$

$$u_6(x,t) = -\frac{\mu^2}{3} - \frac{\mu^2}{2}\cot^2\left(\mu\left(x - \frac{256\mu^6}{3}t\right)\right) - \frac{\mu^2}{2}\tan^2\left(\mu\left(x - \frac{256\mu^6}{3}t\right)\right).$$

3.2. **Solutions by a sinh–cosh rational ansatz.** We change the sinh and cosh functions to their exponential form and then we substitute (3.4) into (3.2). As a result, we obtain a polynomial equation in the variable $\zeta = \exp(\mu\xi)$. Equating the coefficients of the different powers of ζ to zero results in an algebraic system in the variables c, d, k, p, λ and μ. Solving it with the aid of a computer yields following solutions of (1.4) :

- $c = 0, \quad d = -1, \quad k = \frac{\mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_7(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 - \cosh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_8(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 - \cos\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

- $c = 0, \quad d = 1, \quad k = \frac{\mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_9(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 + \cosh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_{10}(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 + \cos\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

- $c = -\sqrt{-1}, \quad d = 0, \quad k = \frac{\mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_{11}(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 - \sqrt{-1}\sinh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_{12}(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 + \sin\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

- $c = \sqrt{-1}, \quad d = 0, \quad k = \frac{\mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_{13}(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 + \sqrt{-1}\sinh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_{14}(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 - \sin\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

- $c = \sqrt{d^2 - 1}, \quad k = \frac{mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_{15}(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 + d\cosh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right) + \sqrt{d^2-1}\sinh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_{16}(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 + d\cos\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right) + \sqrt{1-d^2}\sin\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

- $c = -\sqrt{d^2-1}, \quad k = \frac{\mu^2}{4}, \quad p = -\frac{\mu^2}{24}, \quad \lambda = \frac{\mu^6}{48}$:

$$u_{17}(x,t) = -\frac{\mu^2}{24} + \frac{\mu^2}{4\left(1 + d\cosh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right) - \sqrt{d^2-1}\sinh\left(\mu\left(x + \frac{\mu^6}{48}t\right)\right)\right)}.$$

$$u_{18}(x,t) = \frac{\mu^2}{24} - \frac{\mu^2}{4\left(1 + d\cos\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right) - \sqrt{1-d^2}\sin\left(\mu\left(x - \frac{\mu^6}{48}t\right)\right)\right)}.$$

4. Conclusions

We obtained solutions for the seventh-order KK equation (1.4) by using three distinct methods. These methods are direct and effective and they are applicable to solve other seventh-order KdV equations. We think that the results given here are new in the literature.

References

[1] Y. Pomeau & A. Ramani & B. Grammaticos, *Structural stability of the Korteweg-de Vries solitons under a singular perturbation*, Physica D. 31 (1) (1988) 127–134.

[2] U. Gotkas & W. Hereman, Symbolic computation of conserved densities for systems of nonlinear evolution equations, *J. Symbolic Computation*,1, 1-31, (1999).

[3] J. D. Cole, A quasi-linear parabolic equation occuring in aerodynamics,*Quart. Appl. Math.*, 1951, 9, pages 225-236.

[4] E. Hopf, The partial differential equation $u_t + uu_x = u_{xx}$, *Comm. Pure Appl. Math.*, **3**, 1950, 201-230.